

NON INJECTIVITY OF THE “HAIR” MAP

BERTRAND PATUREAU-MIRAND

ABSTRACT. In [4] an invariant of knots Z^{rat} with values in a space of diagrams with beads has been constructed. When composed with the so called “hair” map H , It gives the Kontsevich integral of the knot. We introduce a new grading on diagrams with beads and use it to show that a non trivial element constructed with Vogel’s zero divisor in the algebra Λ is in the kernel of H . This shows that H is not injective.

INTRODUCTION

The Kontsevich integral Z is the universal rational finite type invariant for knots (see [1]). For a knot K , $Z(K)$ lives in the space of Chinese diagrams isomorphic to $\hat{\mathcal{B}}(*)$ (see section 1.1). Rozansky conjectures ([7]) and Kricker proves ([5]) that Z could be organized into series of “lines” which could be represented by finite $\mathbb{Q}$ -linear combinations of diagrams whose edges were labelled, in an appropriate sense, with rational functions. For knots with trivial Alexander polynomial, this means that $Z(K)$ is in the image of the hair map H (see section 1.3). In [4], It is directly showed that the map Z^{rat} with values in a space of diagrams with beads is an isotopy invariant and Z factor through Z^{rat} . Rozansky, Garoufalidis and Kricker conjecture (see [6] conjecture 3.18) that H could be injective. The theorem 4 shows that this is false.

1. THE HAIR MAP

1.1. Classical diagrams. Let X be a finite set. An X -diagram is an isomorphism class of finite $1 - 3$ -valent graph K with the following data:

- A cyclic ordering on the three oriented edges starting from every trivalent vertices x of K .
- An isomorphism between the set of univalent vertices of K and the set X .

We define $\mathcal{A}(X)$ to be the quotient of the $\mathbb{Q}$ -vector space generated by X -diagram by the relations:

1. The (AS) relations for antisymmetry:

$$\begin{array}{c} \diagup \\ | \\ \diagdown \end{array} + \begin{array}{c} \diagdown \\ | \\ \diagup \end{array} = 0$$

2. The (IHX) relations for three diagrams which differ only in a neighborhood of an edge:

$$\begin{array}{c} \text{---} \text{---} \text{---} \\ | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \text{---} \text{---} \\ | \\ \text{---} \end{array} - \begin{array}{c} \diagup \quad \diagdown \\ | \\ \diagdown \quad \diagup \end{array}$$

These spaces are graduated, the degree of an X -diagram being given by half the total number of vertices.

Set $[n] = \{1, 2, \dots, n\}$ and define F_n to be the subspace of $\mathcal{A}([n])$ generated by

connected diagram with at least one trivalent vertex. The permutation group $\mathfrak{S}(X)$ acts on $\mathcal{A}(X)$ and we will denote by $B(*)$ the coinvariants for this action:

$$B(*) = \bigoplus_{n \in \mathbb{N}} \mathcal{A}([n]) \otimes_{\mathfrak{S}_n} \mathbb{Q}$$

and by $\hat{B}(*)$ its completion for the grading.

Finally Λ will denote Vogel's algebra formed by totally antisymmetric elements of F_3 (for the action of $\mathfrak{S}_3$).

We recall (see [9]) that Λ acts on the modules F_n and that for this action, F_0 and F_2 are free Λ -modules of rank one. Furthermore, the following elements are in Λ :

$$t = \triangle = \frac{1}{2} \bigcirc \quad x_n = \triangle_{n-2}$$

Theorem 1. (Vogel [10]) *The element t is a divisor of zero in Λ .*

Corollary 2. *There exists an element $r \in \Lambda \setminus \{0\}$ such that $t.r = 0$. So one has*

$$r \bigcirc \neq 0 \in F_0 \quad \text{but} \quad r \bigcirc_{1,2,3} = 0 \in F_3$$

Proof: This is just a consequence of the fact that F_0 is a rank one free Λ -module generated by the diagram Θ . The preceding element of F_3 is in fact the product $r \cdot \bigcirc_{1,2,3} = 2tr = 0 \in \Lambda$. $\square$

Remark: Vogel's computation shows that r can be chosen in degree fifteen (in Λ), living in the algebra generated by the x_n and is killed by all the weight systems coming from Lie algebras (but is not killed by the Lie superalgebras $\mathfrak{D}_{2,1,\alpha}$).

1.2. Diagrams with beads. Diagrams with beads have been introduced by Kricker and Garoufalidis (see [5], [4]). A presentation of $\mathcal{B}$ which uses the first cohomology classes of diagrams is already present in [7]. Vogel explains me this point of view for diagrams with beads.

Let G be the multiplicative group $\{b^n, n \in \mathbb{Z}\} \simeq (\mathbb{Z}, +)$ and consider its group algebra $R = \mathbb{Q}G = \mathbb{Q}[b, b^{-1}]$. We denote by $a \mapsto \bar{a}$ the involution of the $\mathbb{Q}$ -algebra R sending b to b^{-1} .

A diagram with beads in R is an $\emptyset$ -diagram with the following supplementary data: The beads which consist of a map $f : E \rightarrow R$ from the set of oriented edges of K such that if $-e$ denotes the same edge than e with opposite orientation, one has $f(-e) = \overline{f(e)}$.

We will represent the beads by some arrows on the edges with label in R . The value of the bead f on e is given by the product of these labels and we won't represent the beads with value 1. So one has the notations:


$$\xrightarrow{f(b)} = \xrightarrow{\overline{f(b)}} \quad \text{and} \quad \xrightarrow{f(b)} \xrightarrow{g(b)} = \xrightarrow{f(b)g(b)}$$

The loop degree of a diagram with beads K is the rank of the first homology group of the underlying graph.

Let $\mathcal{A}^R(\emptyset)$ denotes the quotient of the $\mathbb{Q}$ -vector space generated by diagrams with beads in R by the following relations:

1. (AS)
2. The (IHX) relations are to be consider only near an edge with bead 1.

3. PUSH:



4. Multilinearity:

$$\xrightarrow{\alpha f(b) + \beta g(b)} = \alpha \xrightarrow{f(b)} + \beta \xrightarrow{g(b)}$$

$\mathcal{A}^R(\emptyset)$ is graded by the loop degree:

$$\mathcal{A}^R(\emptyset) = \bigoplus_{n \in \mathbb{N}} \mathcal{A}_n^R(\emptyset)$$

We will prefer an other presentation of $\mathcal{A}^R(\emptyset)$:

- Remark that it is enough to consider diagrams with beads in G and the multilinearity relation can be viewed as a notation.
- Next remark that for a diagrams with beads in G , the map f is the same that a 1-cochain $\tilde{f}$ with value in $\mathbb{Z} \simeq G$ on the underlying simplicial set of K . The elements $\tilde{f}$ are in fact the 1-cocycles because of the condition $f(-e) = \overline{f(e)}$ which means $\tilde{f}(-e) = -\tilde{f}(e)$.

- The “PUSH” relation at a vertex v say that $\tilde{f}$ is only given up to the coboundary of the 0-cochain with value 1 on v and 0 on the other vertices. Hence $\mathcal{A}^R(\emptyset)$ is also the $\mathbb{Q}$ -vector space generated by the pairs (3-valent graph D , $x \in H^1(D, \mathbb{Z})$) quotiented by the relations (AS) and (IHX). With these notations one can describe the (IHX) relations in the following way:

Let K_I , K_H and K_X be three graphs which occur in a (IHX) relation on a edge e . Let $K_\bullet$ be the graph obtained by collapsing the edge e . The map $p_\gamma : K_\gamma \longrightarrow K_\bullet$ induce three cohomology isomorphisms.

If $x \in H^1(K_\bullet, \mathbb{Z})$ then the (IHX) relation at e says that

$$(K_I, p_I^* x) = (K_H, p_H^* x) - (K_X, p_X^* x)$$

is true in $\mathcal{A}^R(\emptyset)$.

1.3. The hair map. The hair map $H : \mathcal{A}^R(\emptyset) \longrightarrow \hat{\mathcal{B}}(*)$ replaces beads by legs (or hair): Just replace a bead t^n by the exponential of n times a leg.

It is well defined (see [4]).

2. GRADING ON DIAGRAMS WITH BEADS

Remark that for a 3-valent graph K , $H^1(K, \mathbb{Z})$ is a free $\mathbb{Z}$ -module. The beads $x \in H^1(K, \mathbb{Z})$ which occur in an (AS) or (IHX) relation are the same up to isomorphisms. We will call p the bead degree of (K, x) if x is p times an indivisible element of $H^1(K, \mathbb{Z})$.

Theorem 3. *The bead degree is well defined in $\mathcal{A}_n^R(\emptyset)$. Thus we have a grading*

$$\mathcal{A}_n^R(\emptyset) = \bigoplus_{p \in \mathbb{N}} \mathcal{A}_{n,p}^R(\emptyset)$$

where $\mathcal{A}_{n,p}^R(\emptyset)$ is the subspace of $\mathcal{A}_n^R(\emptyset)$ generated by diagrams with bead degree p . Furthermore, $\mathcal{A}_{n,0}^R(\emptyset) \simeq \mathcal{A}_n(\emptyset)$ and for $p > 0$, $\mathcal{A}_{n,p}^R(\emptyset) \simeq \mathcal{A}_{n,1}^R(\emptyset)$.

3. A NON TRIVIAL ELEMENT IN THE KERNEL OF H

A diagram of a circle with a horizontal radius line labeled r . The upper arc of the circle is labeled $b-1$ with an arrow pointing clockwise.

$$r \xrightarrow{b-1} r + \frac{1}{2!} r + \frac{1}{3!} r + \dots$$

Remark: The element of theorem 4 has a loop degree seventeen. The hair map is evidently injective on the space of diagrams with bead degree zero. I don't know if it is still true in the other degree.

- [1] D. Bar-Natan - *On the Vassiliev knot invariants*, Topologie 34 n° 2 1995, 423-472.
- [2] D. Bar-Natan, S. Garoufalidis, - *On the Melvin-Morton-Rozansky conjecture*, Invent. Math. 125 (1996) 103-133.
- [3] D. Bar-Natan, S. Garoufalidis, L. Rozansky, D. P. Thurston, - *Wheels, wheeling, and the Kontsevich integral of the unknot*, Hebrew University, Harvard University, University of Illinois at Chicago and University of California at Berkeley preprint, March 1997. q-alg/9703025.
- [4] S. Garoufalidis, A. Kriker - *A rational noncommutative invariant of boundary links* preprint math.GT/0105028, May 2001.
- [5] A. Kriker - *The lines of the Kontsevich integral and Rozansky's rationality conjecture*, Tokyo Institute of Technology preprint, May 2000, arXiv:math.GT/0005284
- [6] T. Ohtsuki - *Problems on Invariants of Knots and 3-Manifold* to appear in Geometry and Topology Monographs.
- [7] L. Rozansky - *Une structure rationnelle sur des fonctions génératrices d'invariants de Vassiliev* Summer school of the University of Grenoble preprint, July 1999.
- [8] L. Rozansky - *A rationality conjecture about Kontsevich integral of knots and its implications to the structure of the colored Jones polynomial*, math.GT/0106097 .
- [9] P. Vogel - *Algebraic structures on modules of diagrams*, preprint to appear Inventiones Mathematicae.
- [10] P. Vogel - *private communication*.

INSTITUT DE MATHÉMATIQUES DE JUSSIEU, EQUIPE “TOPOLOGIE ET GÉOMÉTRIE ALGÈBRIQUES”,
CASE 7012, UNIVERSITÉ PARIS VII, 75251 PARIS CEDEX 05, FRANCE
E-mail address: `patureau@math.jussieu.fr`
URL: `http://www.math.jussieu.fr/~patureau`